



ZYMO RESEARCH

DNA
Purification
MADE SIMPLE
Made Simple™

ZR-96 DNA Clean & Concentrator®-5

Ultra-pure DNA from PCR, enzymatic reactions, and other sources.

Highlights

- Quick, high throughput recovery of ultra-pure DNA from PCR, enzymatic reactions, and other sources.
- DNA can be eluted in as little as 10 μ l and is ideal for DNA ligation, sequencing, labeling, PCR, microarray, transfection, transformation, restriction digestion, etc.

Catalog Numbers:
D4023, D4024



Scan with your smart-phone camera to
view the online protocol/video.



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682

Table of Contents

Product Contents	01
Specifications	02
Product Description	03
Formats	04
Applications	04
Protocol	05
Buffer Preparation.....	05
Sample Processing.....	05
Appendix	06
ssDNA Purification.....	06
M13 phage ssDNA Purification	06
Isolated DNA stored in DNA/RNA Shield ..	07
RNase A Treatment	07
Troubleshooting	08
Ordering Information	09
Complete Your Cloning Workflow	10
Notes	11
Guarantee	13

Product Contents

ZR-96 DNA Clean & Concentrator®-5	D4023 (2 x 96 Preps.)	D4024 (4 x 96 Preps.)	Storage Temperature
DNA Binding Buffer	100 ml	2 x 100 ml	Room Temp.
DNA Wash Buffer ¹	24 ml	48 ml	Room Temp.
DNA Elution Buffer	10 ml	16 ml	Room Temp.
Zymo-Spin™ I-96 Plate	2	4	Room Temp.
Collection Plate	2	4	Room Temp.
Elution Plate	2	4	Room Temp.
Instruction Manual	1	1	-


¹ Ethanol must be added prior to use as indicated on the **DNA Wash Buffer** label.

Specifications


- **DNA Purity** – High-quality DNA ($A_{(260/280)} \geq 1.8$) ideal for ligation, sequencing, labeling, PCR, microarray, transfection, transformation, and restriction digestion procedures.
- **DNA Size Limits** – From ~50 bp to 23 kb.
- **DNA Recovery** – Typically, up to 5 μg total DNA per well can be eluted into as little as 10 μl of low salt **DNA Elution Buffer** or water. For DNA 50 bp to 10 kb, the recovery is 70-90%. For DNA 11 kb to 23 kb, the recovery is 50-70%.
- **Sample Sources** – DNA from enzymatic reactions (e.g., PCR, restriction endonuclease digestions), plasmid preparations, and impure preparations. Suitable for isolated DNA stored in DNA/RNA Shield (page 7).
- **Product Detergent Tolerance** – $\leq 5\%$ Triton X-100, $\leq 5\%$ Tween-20, $\leq 5\%$ Sarkosyl, $\leq 0.1\%$ SDS.

Product Description


The **ZR-96 DNA Clean & Concentrator[®]-5 (ZR-96 DCC[®]-5)** provides a hassle-free method for the rapid, high throughput purification and concentration of high-quality DNA from PCR, endonuclease digestions, cell lysates and other impure DNA preparations. It can also be used for post-RT cDNA clean-up and purification of sequencing-ready DNA from M13 phage. Simply add the specially formulated **DNA Binding Buffer** to your sample and transfer to the wells of the supplied **Zymo-Spin[™] I-96 Plate**. There is no need for organic denaturants or chloroform. Instead, the product features *Fast-Spin* technology to yield DNA that is free of salts and contaminants in just minutes. DNA purified using the **ZR-96 DCC[®]-5** is suitable for nucleotide sequencing, microarray analysis, PCR, and restriction endonuclease digestion procedures.



ZR-96 DCC[®]-5 procedure.



Clean & Concentrated DNA. DNA samples, such as the PCR products shown here, can be efficiently purified and concentrated using the DNA Clean & Concentrator[®]-5.



Pure and Reliable Recovery with the DCC[®]-5. Shown here is the recovery of 1 µg of 100 bp marker DNA eluted into 10 µl of water analyzed using a NanoDrop[®] spectrophotometer. The **DNA Clean & Concentrator[®]-5** consistently recovers > 90 % of input DNA.

Formats

	DCC™-5	DCC™-25	DCC™-100	DCC™-500	Genomic DCC™	ZR-96 DCC™-5
						
Name	Zymo-Spin™ I & IC	Zymo-Spin™ II & IIC	Zymo-Spin™ V	Zymo-Spin™ VI	Zymo-Spin™ IC-XL	Zymo-Spin™ I-96
Capacity	5 µg/ prep.	25 µg/ prep.	100 µg/ prep.	500 µg/ prep.	10 µg/ prep.	5 µg/ prep.
Elution Vol.	≥ 6 µl	≥ 25 µl	≥ 150 µl	≥ 2 ml	≥ 10 µl	≥ 10 µl
Cat. Nos.	D4003, D4013	D4005, D4033	D4029, D4030	D4031, D4032	D4010, D4011	D4023, D4024

Applications

Post-PCR DNA Clean-up	Efficient desalting of DNA with the removal of DNA polymerases, primers, and free dNTPs.
DNA Clean-up From Enzymatic Reactions	Efficient desalting of DNA with the removal of modifying enzymes, RNA polymerases, ligases, kinases, nucleases, phosphatases, endonucleases, etc.
Post-Reverse Transcription (RT) & cDNA Clean-up	Efficiently purifies DNA following RT, either as a DNA/RNA complex or as single stranded cDNA following chemical hydrolysis of the RNA template.
Plasmid DNA Clean-up	Efficiently purifies plasmid DNA from “home-made” preparations of cell free lysates or from commercial kits. Plasmid DNA purified and concentrated using the DCC® has proven an excellent substrate for high quality DNA sequencing.
Isotope and Dye Removal	Efficiently removes unincorporated fluorescent (<i>i.e.</i> , AMCA, FITC, BIO, DIG, Cy3, Cy5, FAM, etc.) and radiolabeled dNTP derivatives from DNA following <i>in vitro</i> labeling reactions.
Purification of M13 ssDNA	The DCC® can be used for the rapid isolation of single stranded M13 phage DNA directly from phage-infected <i>E. coli</i> culture supernatant.

✓ For purification of short DNA or RNA oligonucleotides ≥ 16 nt, use the **Oligo Clean & Concentrator™ (D4060, D4061)**.

✓ For ChIP (Chromatin Immunoprecipitation) sample cleanup, use the **ChIP DNA Clean & Concentrator® (D5201, D5205)** for high quality DNA from any step in a standard ChIP protocol.

✓ For post-cycle sequencing samples, use the **ZR Sequencing DNA Clean-up Kit™ (D4050, D4051)** for dye blob elimination.

✓ For samples containing PCR inhibitors, use the **OneStep™ PCR Inhibitor Removal Kit (D6030, D6035)**.

Protocol

Buffer Preparation

- ✓ **Before starting:** Add 96 ml 100% ethanol (104 ml 95% ethanol) to the 24 ml **DNA Wash Buffer** concentrate. Add 192 ml 100% ethanol (208 ml 95% ethanol) to the 48 ml **DNA Wash Buffer** concentrate.

Sample Processing

All centrifugation steps should be performed between 3,000 - 5,000 x g.

1. Add 2-7 volumes of **DNA Binding Buffer** to each volume of DNA sample (see table below)¹. Mix briefly by vortexing.

Application	DNA Binding Buffer : Sample	Example
Plasmid, genomic DNA (>2 kb) ²	2 : 1	200 µl : 100 µl
PCR product, DNA fragment	5 : 1	500 µl : 100 µl
ssDNA ³ (e.g. cDNA, M13 phage)	7 : 1	700 µl : 100 µl

2. Transfer sample mixtures to the wells of a **Zymo-Spin™ I-96 Plate**⁴ mounted on a **Collection Plate**.
3. Centrifuge for 5 minutes until sample mixtures have been completely filtered. Discard the flow-through.
4. Add 300 µl **DNA Wash Buffer** to each well of the **Zymo-Spin™ I-96 Plate**. Centrifuge for 5 minutes. Repeat wash step, but centrifuge for 15 minutes. (Alternatively, one wash can be performed using 600 µl **DNA Wash Buffer**).
5. Add ≥ 10 µl **DNA Elution Buffer**⁵ or water⁶ directly to the column matrix in each well. Transfer the **Zymo-Spin™ I-96 Plate** onto an **Elution Plate** and centrifuge for 3 minutes to elute the DNA. Ultra-pure DNA is now ready for use.

¹ Add a minimum of 100 µl of **DNA Binding Buffer** to all samples ≤ 50 µl.

² For efficient recovery of DNA >20 kb, use the ZR-96 **Genomic DNA Clean & Concentrator-5 (D4066, D4067)**.

³ For ssDNA purification, see **Appendix A** on page 5.

⁴ The capacity of each well of the **Zymo-Spin™ I-96 Plate** is approximately 1.1 ml. The capacity of each well of the **Collection Plate** is approximately 800 µl. Therefore, it may be necessary to load and spin the plate multiple times if a sample has a volume larger than 800 µl.

⁵ **DNA Elution Buffer:** 10mM Tris-HCl, pH 8.5, 0.1mM EDTA.

⁶ Elution of DNA from the column is dependent on pH and temperature. If water is used, make sure the pH is > 6.0. Waiting 1 minute prior to elution may improve the yield of larger (> 6 kb) DNA. For even larger DNA (> 10 kb), the total yield may be improved by eluting the DNA with 60-70°C **DNA Elution Buffer**.

Appendix

cDNA clean-up

The **DCC**[®] kit can be used to effectively clean and concentrate cDNA (> 500 nt) following reverse transcription (RT) in the presence/absence of fluorescent dyes. Unincorporated free nucleotides and fluorescent derivatives are efficiently removed using the **DCC**[®], and the recovered cDNA may be used directly for microarray analysis, second-strand cDNA synthesis, or indirect labeling with a fluorescent dye such as NHS ester Cy3 or Cy5.

For clean-up of short cDNAs or ESTs (≥ 16 nt), we recommend the **Oligo Clean & Concentrator** (Cat. Nos. **D4060**, **D4061**).

Hydrolysis

1. Add 10 μ l 0.5 M EDTA and 10 μ l 1 N NaOH to 50 μ l of RT reaction.

The volumes of EDTA and NaOH should be scaled proportionally depending on the starting volume of the RT reaction.

2. Incubate at 65°C for 15 minutes.

Clean-up

1. Add 490 μ l (7 volumes) of **DNA Binding Buffer** to the hydrolysis reaction above. Mix well.

Neutralization (pH) following RNA hydrolysis is not necessary as the **DNA Binding Buffer** will effectively neutralize the NaOH added to the reaction.

2. Continue with Step 2 of the Sample Processing Protocol on page 5.

M13 phage ssDNA purification

1. Centrifuge phage-infected bacterial culture at 8,000 x g for 1 minute.
2. Transfer 100 μ l of phage-containing supernatant to a 1.5 ml microcentrifuge tube and add 700 μ l (7 volumes) of **DNA Binding Buffer**. Mix briefly by vortexing.

Increased supernatant volumes may be processed by proportionally increasing the amount of **DNA Binding Buffer** added to the sample.

3. Continue with Step 2 of the Sample Processing Protocol on page 5.

Isolated DNA stored in DNA/RNA Shield

*For previously isolated/purified DNA stored in **DNA/RNA Shield**, use the following protocol to recover ultra-pure DNA, ready for downstream applications.*

1. If frozen, thaw samples¹ at room temperature (20-30°C).
2. Add an equal volume of ethanol (95-100%) to the sample and mix well.
3. Continue with Step 2 of the Sample Processing Protocol on page 5.

RNase A Treatment

Dissolve RNase A (E1008-30), sold separately, in DNase/RNase-free water or TE to a stock concentration of 10 mg/ml.

1. Add enough 10 mg/ml RNase A to the sample for a final concentration of 10-100 µg/mL and mix well.
2. Add Incubate at room temperature for 15 minutes.
3. Continue with step 1 of the Sample Processing protocol on page 5.

¹Adjust the sample volume to 50 µl (minimum) with **DNA/RNA Shield**.

Troubleshooting

Problem	Possible Causes and Suggested Solutions
Low Recovery	<p>Improperly Prepared/Stored DNA Wash Buffer. Make sure ethanol has been added to the DNA Wash Buffer concentrate. Cap the bottle tightly to prevent evaporation over time.</p> <p>Addition of DNA Elution Buffer. Add elution buffer directly to the column matrix, not to the walls of the column. Elution buffer requires contact with the matrix for at least 1 minute for large DNA ≥ 10 kb.</p> <p>Incomplete Elution. DNA elution is dependent on pH, temperature, and time. For large genomic DNA (≥ 50 kb), apply heated elution buffer (60-70 °C) to the column and incubate for several minutes prior to elution. Sequential elutions may be performed for quantitatively higher recovery but lower final DNA concentration. This is recommended for DNA ≥ 10 kb.</p>
Low A_{260}/A_{230} ratio	<p>Column tip contaminated. When removing the column from the collection tube, be careful that the tip of the column does not come into contact with the flowthrough. Trace amounts of salt from the flowthrough can contaminate a sample resulting in a low A_{260}/A_{230} ratio. Ethanol contamination from the flowthrough can also interfere with DNA elution. Zymo-Spin™ columns are designed for complete elution with no buffer retention or carryover.</p>
Following Clean-up with DCC®, Multiple Bands Appear in an Agarose Gel	<p>Acidification of DNA Loading Dye. Most loading dyes do not contain EDTA and will acidify ($\text{pH} \leq 4$) over time due to some microbial growth. This low pH is enough to cause DNA degradation. Therefore, if water is used to elute the DNA, 6X Loading Dye containing 1 mM EDTA is recommended.</p>

Ordering Information

Product Description	Catalog No.	Size
DNA Clean & Concentrator[®]-5 (for purification of up to 5 µg DNA per prep.)	D4003T (uncapped)	10 Preps.
	D4003 (uncapped)	50 Preps.
	D4004 (uncapped)	200 Preps.
	D4013 (capped)	50 Preps.
ZR-96 DNA Clean & Concentrator[®]-5 (for 96-well purification of up to 5 µg DNA per well)	D4023	2 x 96 Preps.
	D4024	4 x 96 Preps.
DNA Clean & Concentrator[®]-25 (for purification of up to 25 µg DNA per prep.)	D4005 (uncapped)	50 Preps.
	D4006 (uncapped)	200 Preps.
	D4033 (capped)	50 Preps.
	D4034 (capped)	200 Preps.
DNA Clean & Concentrator[®]-100 (for purification of up to 100 µg DNA per prep.)	D4029	25 Preps.
	D4030	50 Preps.
DNA Clean & Concentrator[®]-500 (for purification of up to 500 µg DNA per prep.)	D4031	10 Preps.
	D4032	20 Preps.

Individual Kit Components	Catalog No.	Amount
DNA Binding Buffer	D4003-1-L	50 ml
	D4004-1-L	100 ml
DNA Wash Buffer (concentrate)	D4003-2-24	24 ml
	D4003-2-48	48 ml
DNA Elution Buffer	D3004-4-10	10 ml
	D3004-4-16	16 ml
Zymo-Spin™ I-96 Plate	C2004-2	2 Plates
Collection Plate	C2002	2 Plates
Elution Plate	C2003	2 Plates

Complete Your Cloning Workflow

✓ Transfection-grade plasmid DNA from a miniprep

ZymoPURE™ Plasmid Miniprep	Size	Catalog No.
ZymoPURE™ Plasmid Miniprep Kit	10 Preps. 50 Preps. 100 Preps. 400 Preps. 800 Preps.	D4208T D4209 D4210 D4211 D4212

✓ 20 Minute Endotoxin-Free Midi & Maxipreps

ZymoPURE™ II Plasmid Prep Kits	Size	Catalog No.
ZymoPURE™ II Plasmid Midiprep Kit	25 Preps. 50 Preps.	D4200 D4201
ZymoPURE™ II Plasmid Maxiprep Kit	10 Preps. 20 Preps.	D4202 D4203
ZymoPURE™ II Plasmid Gigaprep Kit	5 Preps.	D4204

✓ Simple 20 second High Efficiency Transformations

Mix & Go! Competent Cells	Size	Catalog No.
DH5α	10 x 100 µl aliquots 96 x 50 µl aliquots 96 x 50 µl aliquots PCR Plate	T3007 T3009 T3010
Zymo10B	10 x 100 µl aliquots 96 x 50 µl aliquots	T3019 T3020
JM109	10 x 100 µl aliquots 96 x 50 µl aliquots	T3003 T3005
HB101	10 x 100 µl aliquots 96 x 50 µl aliquots	T3011 T3013
TG1	10 x 100 µl aliquots	T3017

✓ Recover ultra-pure highly concentrated DNA from PCR & other sources

DNA Clean & Concentrator™	Size	Catalog No.
DNA Clean & Concentrator™ -5	50 Preps. 200 Preps.	D4003 D4004
ZR-96 DNA Clean-Up Kit™	2 x 96 Preps. 4 x 96 Preps.	D4017 D4018

✓ Rapid extraction of ultra-pure DNA from agarose gels

Zymoclean Gel DNA Recovery™	Size	Catalog No.
Zymoclean™ Gel DNA Recovery Kit	50 Preps. 200 Preps.	D4001 D4002



**100% satisfaction guarantee on all Zymo Research products,
or your money back.**

Zymo Research is committed to simplifying your research with quality products and services. If you are dissatisfied with this product for any reason, please call 1(888) 882-9682.

Integrity of kit components is guaranteed for up to one year from date of purchase.
Reagents are routinely tested on a lot-to-lot basis to ensure they provide the highest performance and reliability.

This product is for research use only and should only be used by trained professionals. It is not for use in diagnostic procedures. Some reagents included with this kit are irritants. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

TM Trademarks of Zymo Research Corporation
NanoDrop® is a registered trademark of NanoDrop Technologies, Inc.



ZYMO RESEARCH

The **BEAUTY** of **SCIENCE** is to Make Things **SIMPLE**®



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682