



ZYMO RESEARCH

DNA
Purification
MADE SIMPLE
Made Simple™

ZymoPURE™ II Plasmid Maxiprep Kit

Rapid purification of endotoxin-free plasmid DNA from up to 150 ml of overnight *E. coli* culture.

Highlights

- Perform plasmid maxipreps in only 16 minutes using a simple spin-column protocol.
- Purify up to 3.0 mg of highly concentrated plasmid DNA directly from a spin-column.
- Eluted plasmid DNA is Endotoxin-free and Transfection-Ready.

Catalog Numbers:
D4202, D4203



Scan with your smart-phone camera to
view the online protocol/video.



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682

Table of Contents

Product Contents	01
Specifications	02
Product Description	03
Procedure Overview	04
Protocol	05
Buffer Preparation.....	05
Plasmid DNA Purification.....	05
Appendices	08
Low-Copy Number Plasmid Protocol	08
Gram-Positive Bacteria Protocol	09
Adjusting Volume of Binding Buffer	10
Troubleshooting	11
Ordering Information	13
Complete Your Cloning Workflow	14
Notes	15
Guarantee	17

Product Contents

ZymoPURE™ II Plasmid Maxiprep Kit	D4202 (10 prep)	D4203 (20 prep)	Storage Temperature
ZymoPURE™ P1 ¹ (Red)	150 ml	150 ml (2x)	4°C
ZymoPURE™ P2 ^{2,3} (Blue)	150 ml	150 ml (2x)	Room Temp.
ZymoPURE™ P3 (Yellow)	150 ml	150 ml (2x)	Room Temp.
ZymoPURE™ Binding Buffer ³	150 ml	150 ml (2x)	Room Temp.
ZymoPURE™ Wash 1	55 ml (2x)	55 ml (4x)	Room Temp.
ZymoPURE™ Wash 2 (concentrate) ⁴	23 ml (2x)	23 ml (4x)	Room Temp.
ZymoPURE™ Elution Buffer	6 ml	12 ml	Room Temp.
Zymo-Spin™ V-PX Column Assemblies ⁵	10 pcs	20 pcs	Room Temp.
ZymoPURE™ Syringe Filter-X	10 pcs	20 pcs	Room Temp.
ZymoPURE™ Syringe Plungers	10 pcs	20 pcs	Room Temp.
EndoZero™ Spin-Columns	10 pcs	20 pcs	Room Temp.
Collection Tubes	10 pcs	20 pcs	Room Temp.
Instruction Manual	1 pc	1 pc	-

¹ ZymoPURE™ P1 contains RNase A (100 µg/ml) and is stable at room temperature without loss in RNase activity, however, for long-term storage the product should be stored at 4-8° C.

² Caution: ZymoPURE™ P2 Buffer contains NaOH. Please use proper safety precautions.

³ The ZymoPURE™ P2 and ZymoPURE™ Binding Buffer may have precipitated. If this occurs, dissolve the precipitate by incubating the bottles at 30-37 °C for 10-20 minutes and mix by inversion. Do not microwave!

⁴ ZymoPURE™ Wash 2 included with D4202 and D4203 are supplied as a concentrate and require the addition of ethanol prior to use. See Buffer Preparation (page 5) for instructions.

⁵ The Zymo-Spin™ V-PX, 15 ml Reservoir-X and 50 ml Reservoir are pre-assembled as a single unit.

Specifications

- **DNA Purity** – Eluted DNA is ultrapure, endotoxin-free, and well suited for transfection, transformation, lentivirus production, adenovirus production, AAV production, CRISPR, genome editing, *in vivo* studies, sequencing, restriction endonuclease digestion, *in vitro* transcription/translation, PCR, and other sensitive applications.
 - Typical Abs260/280 ≥ 1.8 and Abs260/230 ≥ 2.0
 - Endotoxin levels: ≤ 1 EU/ μ g of plasmid DNA using the Standard Protocol
 - ≤ 0.025 EU/ μ g of plasmid DNA with optional EndoZero™ II Spin-Column
- **Plasmid DNA Yield** – Up to 3.0 mg per preparation. Actual yield is dependent on the plasmid copy number, culture growth conditions, and strain of *E. coli* utilized. Typical yields from 150 ml of overnight culture grown in LB for high-copy plasmids are 600 – 1,200 μ g.
- **Plasmid DNA Size** – Up to ~200 kb
- **Recovery Volume** – ≥ 300 μ l of ZymoPURE™ Elution Buffer or DNase-free water
- **Processing Time** – ≤ 18 min
- **Required Equipment** – Microcentrifuge and vacuum/vacuum manifold (recommended) or swinging-bucket centrifuge.

Product Description

The **ZymoPURE™ II Plasmid Maxiprep Kit** features a simple spin-column based method for the purification of up to 3 mg of transfection grade plasmid DNA in less than 18 minutes. The eluted plasmid DNA is endotoxin-free and ready for immediate use in the most sensitive applications. The unique ZymoPURE™ methodology removes the need for slow gravity flow anion-exchange columns, alcohol precipitation, lengthy endotoxin removal incubations, and time-consuming centrifugation steps.


ZymoPURE™ technology uses a modified alkaline lysis method and features our patented binding chemistry and EZ-Flow™ spin-column design, which enables the highest DNA binding capacity and rapid loading of the lysate and wash buffer, resulting in the fastest purification of highly concentrated (up to 6 mg/ml) plasmid DNA directly from a spin-column. Coupling ZymoPURE™ with the innovative **EndoZero™ Spin-Columns**, to eliminate residual endotoxins, achieves endotoxin-free plasmid DNA (≤ 0.025 EU/ μ g of plasmid DNA), making it suitable for transfection, recombinant virus production, lentivirus production, genome editing, *in vivo* studies, sequencing, restriction endonuclease digestion, *in vitro* transcription/translation, PCR, transformation, and other sensitive applications.

As an added convenience, the **ZymoPURE™ II Plasmid Maxiprep Kit** contains colored buffers that permit error-free visualization and identification of complete bacterial cell lysis and neutralization. Syringe filters are also included for rapid clearing of the lysate and the unique spin-column design allows the binding step to be performed using a vacuum or centrifuge.


Rapid Purification of Highly Concentrated Endotoxin-Free Plasmid DNA



Up to 6x More Concentrated Plasmid



Up to 3x More Plasmid DNA Yield



Yield and concentration for plasmid DNA isolated using the ZymoPURE™ II Maxiprep kit compared to two endotoxin-free kits from Supplier Q and Supplier MN. Plasmid DNA (pGL3®) was isolated from 150 ml of JM109 *E. coli* culture grown overnight following the manufacturer's suggested protocol (in duplicate). One (1) μ l of eluted plasmid DNA was visualized post agarose gel electrophoresis. M, ZR 1 kb DNA Marker (Zymo Research).

Procedure Overview



Bacterial cells are resuspended in **ZymoPURE™ P1** (red).



The solution will turn dark purple and viscous following the addition of **ZymoPURE™ P2** (blue) indicating bacterial lysis is complete.



The solution will turn yellow and a precipitate will form after adding **ZymoPURE™ P3** (yellow) indicating neutralization is complete.



The neutralized lysate is loaded into the **ZymoPURE™ Syringe Filter-X** and clarified into a new 50 ml conical tube.



ZymoPURE™ Binding Buffer is added to the cleared lysate and mixed thoroughly.



The mixture is loaded into the **Zymo-Spin™ V-PX Column** using a vacuum manifold.



The **50 ml Reservoir** is removed and the **Zymo-Spin™ V-PX Column** is washed using a vacuum manifold.



Ultra-pure plasmid DNA is eluted from the **Zymo-Spin™ V-PX Column** using a microcentrifuge.



The eluted plasmid DNA is passed through the **EndoZero™ Column** using a microcentrifuge.

Protocol

Buffer Preparation:

- ✓ Add 88 ml of 95% ethanol to the **23 ml ZymoPURE™ Wash 2 (Concentrate)** before use.
- ✓ The **ZymoPURE™ P2** and **ZymoPURE™ Binding Buffer** may have precipitated. If this occurs, dissolve the precipitate by incubating the bottles at 30-37 °C for 10-20 minutes and mix by inversion. Do not microwave!

Before Starting:

- ✓ Centrifuge up to 150 ml of bacterial culture at $\geq 3,400 \times g$ for 10 minutes to pellet the cells. Discard supernatant.

Plasmid DNA Purification

The following procedure should be performed at room temperature (15-30°C).

1. Add 14 ml of **ZymoPURE™ P1 (Red)** to the bacterial cell pellet and resuspend completely by vortexing or pipetting.
2. Add 14 ml of **ZymoPURE™ P2 (Blue)** and immediately mix by gently inverting the tube 6 times. Do not vortex! Let sit at room temperature for 2-3 minutes¹. *Cells are completely lysed when the solution appears clear, purple, and viscous.*
3. Add 14 ml of **ZymoPURE™ P3 (Yellow)** and mix gently but thoroughly by inversion. Do not vortex! Invert the tube an additional 5 times after the sample turns completely yellow. *The sample will turn yellow when the neutralization is complete, and a yellowish precipitate will form.*
4. Ensure the plug is attached to the Luer Lock at the bottom of the **ZymoPURE™ Syringe Filter-X**. Place the syringe filter upright in a tube rack and load the lysate in the **ZymoPURE™ Syringe Filter-X²** and wait 5-8 minutes for the precipitate to float to the top.
5. Remove the Luer Lock plug from the bottom of the syringe and place it into a clean 50 ml conical tube. Place the plunger in the syringe and push the solution through the **ZymoPURE™ Syringe Filter-X** in one continuous motion until approximately 35 ml of cleared lysate is recovered. Save the cleared lysate!

The volume of lysate recovered from the syringe filter is critical for optimal plasmid DNA binding to the spin-column. Please refer to page 10 in the appendix regarding the adjustment of the volume of ZymoPURE™ Binding Buffer used in step 6 if the clarified lysate volume is below approximately 35 ml.

6. Add 14 ml of **ZymoPURE™ Binding Buffer** to the cleared lysate from step 5 and mix thoroughly by inverting the tube 8 times.

To continue processing the lysate using the recommended vacuum protocol, proceed to the next page. If a vacuum is not available, proceed to page 7 for an alternative centrifugation method.

¹ Do not allow the lysis reaction to proceed for more than 3 minutes. Excessive lysis can result in denatured plasmid DNA.

² If the precipitate has formed a homogenous layer at the surface of the neutralized lysate, then invert the tube 3-4 times prior to loading the lysate into the **ZymoPURE™ Syringe Filter-X**.

Vacuum Protocol:

This product is compatible with any conventional vacuum-based manifold. The vacuum pump should be a single or double-staged unit capable of producing up to 400 mm Hg pressure at the vacuum manifold¹.

7. Ensure the connections of the **Zymo-Spin™ V-PX Column Assembly** are finger-tight and place onto a vacuum manifold. (If vacuum is not available, see page 7 for the centrifugation protocol.)
8. With the vacuum off, add the entire mixture from step 6 into the Zymo-Spin™ V-PX Column Assembly, and then turn on the vacuum¹ until all of the liquid has passed completely through the column.
9. Remove and discard the **50 ml Reservoir** from the top of the Zymo-Spin™ V-PX Column Assembly.
10. With the vacuum off, add 10 ml of **ZymoPURE™ Wash 1** to the Zymo-Spin™ V-PX Column Assembly. Turn on the vacuum until all of the liquid has passed completely through the column.
11. With the vacuum off, add 10 ml of **ZymoPURE™ Wash 2** to the Zymo-Spin™ V-PX Column Assembly. Turn on the vacuum until all of the liquid has passed completely through the column. Repeat this wash step.
12. Remove and discard the 15 ml Reservoir-X and place the **Zymo-Spin™ V-PX Column** in a **Collection Tube**. Centrifuge at $\geq 16,000 \times g$ for 1 minute, in a microcentrifuge, to remove any residual wash buffer.
13. Transfer the column into a clean 1.5 ml microcentrifuge tube and add 400 μ l of **ZymoPURE™ Elution Buffer**^{2,3,4} directly to the column matrix. Wait 2 minutes, and then centrifuge at $\geq 16,000 \times g$ for 1 minute in a microcentrifuge.
14. *Optional:* For removal of residual endotoxins^{5,6}, place the **EndoZero™ Spin-Column** in a clean 1.5 ml microcentrifuge tube. Add the entire eluate from step 13 into the EndoZero™ Spin-Column, wait 2 minutes, and then centrifuge at $10,000 \times g$ for 1 minute in a microcentrifuge. Store the eluted plasmid DNA at $\leq -20^{\circ}\text{C}$.

¹ To achieve optimal performance, the vacuum pump should be able to apply at least 400 mm Hg pressure. If less pressure is applied, centrifuge the column prior to washing to remove any residual lysate remaining in the matrix.

² The **ZymoPURE™ Elution Buffer** contains 10 mM Tris-HCl, pH 8.5, 0.1 mM EDTA. If required, pure water can also be used to elute the DNA.

³ The DNA yield can be increased by pre-warming the **ZymoPURE™ Elution Buffer** to 50 °C and/or increasing the incubation period up to 10 minutes prior to centrifugation.

⁴ For low-copy number plasmids or if higher concentration is desired, the plasmid DNA can be eluted in as little as 300 μ l.

⁵ This optional step will reduce endotoxin levels from ≤ 1 EU/ μ g of plasmid DNA to ≤ 0.025 EU/ μ g of plasmid DNA.

⁶ Due to the **EndoZero™ Spin-Column** chemistry, some plasmid DNA will be lost during this step. The percent of plasmid DNA loss will be dependent on the amount of plasmid DNA that is going through the spin-column. The plasmid DNA loss is generally not significant for moderate and high-copy number plasmids. However, it can be significant for low-copy number plasmids or lower yielding preps.

Centrifugation Protocol:

Perform steps 1-6 as indicated in the general protocol on page 5 and continue with the protocol below using a swinging-bucket centrifuge.

7. Remove the **50 ml Reservoir** from the top of the **Zymo-Spin™ V-PX Column Assembly**. Ensure the connection between the **15 ml Reservoir-X** and **Zymo-Spin™ V-PX Column** is finger-tight and place the assembly into a 50 ml conical tube.
8. Add 10 ml of the mixture from step 6 into the **15 ml Reservoir-X/Zymo-Spin™ V-PX Column Assembly**, and centrifuge at 500 x g for 2 minutes. Empty the 50 ml conical tube and repeat this step until the entire mixture has passed through the column.
9. Add 10 ml of **ZymoPURE™ Wash 1** to the Zymo-Spin™ V-PX Column Assembly and centrifuge the column at 500 x g for 2 minutes. Discard the flow through.
10. Add 10 ml of **ZymoPURE™ Wash 2** to the Zymo-Spin™ V-PX Column Assembly and centrifuge at 500 x g for 2 minutes. Discard the flow through and repeat this wash step.
11. Remove and discard the **15 ml Reservoir-X** from the **Zymo-Spin™ V-PX Column**. Place the Zymo-Spin™ V-PX Column in a **Collection Tube** and centrifuge at $\geq 16,000 \times g$ for 1 minute, in a microcentrifuge, to remove any residual wash buffer.
12. Transfer the Zymo-Spin™ V-PX Column into a clean 1.5 ml tube and add 400 μ l of **ZymoPURE™ Elution Buffer**^{1,2,3} directly to the column matrix. Incubate at room temperature for 2 minutes, and then centrifuge at $\geq 16,000 \times g$ for 1 minute in a microcentrifuge.
13. *Optional:* For removal of residual endotoxins^{4,5}, place the **EndoZero™ Spin-Column** in a clean 1.5 ml microcentrifuge tube. Add the entire eluate from step 12 into the EndoZero™ Spin-Column, wait 2 minutes, and then centrifuge at 10,000 x g for 1 minute in a microcentrifuge. Store the eluted plasmid DNA at $\leq -20^\circ\text{C}$.

¹ The **ZymoPURE™ Elution Buffer** contains 10 mM Tris-HCl, pH 8.5, 0.1 mM EDTA. If required, pure water can also be used to elute the DNA.

² The DNA yield can be increased by pre-warming the **ZymoPURE™ Elution Buffer** to 50 °C and/or increasing the incubation period up to 10 minutes prior to centrifugation.

³ For low-copy number plasmids or if higher concentration is desired, the plasmid DNA can be eluted in as little as 300 μ l.

⁴ This optional step will reduce endotoxin levels from ≤ 1 EU/ μ g of plasmid DNA to ≤ 0.025 EU/ μ g of plasmid DNA.

⁵ Due to the **EndoZero™ Spin-Column** chemistry, some plasmid DNA will be lost during this step. The percent of plasmid DNA loss will be dependent on the amount of plasmid DNA that is going through the spin-column. The plasmid DNA loss is generally not significant for moderate and high-copy number plasmids. However, it can be significant for low-copy number plasmids or lower yielding preps.

Appendices

Low-Copy Number Protocol

When working with low-copy number plasmid DNA, it is possible to increase plasmid DNA yield by processing up to 300 ml of overnight culture grown in LB using the protocol below.

Before Starting:

- ✓ Centrifuge up to 300 ml of bacterial culture at $\geq 3,400 \times g$ for 10 minutes to pellet the cells. Discard supernatant.

Plasmid DNA Purification

The following procedure should be performed at room temperature (15-30°C).

1. Add 14 ml of **ZymoPURE™ P1 (Red)** to the bacterial cell pellet and resuspend completely by vortexing or pipetting.
2. Add 14 ml of **ZymoPURE™ P2 (Blue)** and immediately mix by gently inverting the tube 8-10 times. Do not vortex! Let sit at room temperature for 5 minutes. *Cells are completely lysed when the solution appears clear, purple, and viscous.*
3. Add 14 ml of **ZymoPURE™ P3 (Yellow)** and mix gently but thoroughly by inversion. Do not vortex! Invert the tube an additional 5 times after the sample turns completely yellow. *The sample will turn yellow when the neutralization is complete, and a yellowish precipitate will form.*
4. Centrifuge the neutralized lysate for 10 minutes at $\geq 3,400 \times g$.
5. Remove the Luer Lock plug from the bottom of the **ZymoPURE™ Syringe Filter-X** and place it into a clean 50 ml conical tube. Place the plunger in the syringe and push the solution through the ZymoPURE™ Syringe Filter-X in one continuous motion until approximately 35 ml of cleared lysate is recovered. Save the cleared lysate!

The volume of lysate recovered from the syringe filter is critical for optimal plasmid DNA binding to the spin-column. Please refer to page 10 in the appendix regarding the adjustment of the volume of ZymoPURE™ Binding Buffer used in step 6 if the clarified lysate volume is below approximately 35 ml.

6. Add 14 ml of **ZymoPURE™ Binding Buffer** to the cleared lysate from step 5 and mix thoroughly by inverting the capped tube 8 times.

To continue processing the lysate using the recommended vacuum protocol, proceed to page 6. If a vacuum is not available, proceed to page 7 for an alternative centrifugation method.

Gram-Positive Bacteria Protocol

It is possible to isolate plasmid DNA from Gram-Positive species with the ZymoPURE™ II Maxiprep Kit. However, the cell walls of the bacteria must be digested with a lytic enzyme prior to alkaline lysis. The protocol below is for Gram-Positive strains that are sensitive to the lytic enzyme Lysozyme.

Plasmid DNA Purification

The following procedure should be performed at room temperature (15-30°C).

1. Add 14 ml of **ZymoPURE™ P1 (Red)** containing lysozyme¹ at a final concentration of 1 mg/ml to the bacterial cell pellet and resuspend completely by vortexing or pipetting. Incubate the resuspended cell pellet at 37°C for 15-60 minutes².
2. Add 14 ml of **ZymoPURE™ P2 (Blue)** and immediately mix by gently inverting the tube 6 times. Do not vortex! Let sit at room temperature for 2-3 minutes³. *Cells are completely lysed when the solution appears clear, purple, and viscous.*
3. Add 14 ml of **ZymoPURE™ P3 (Yellow)** and mix thoroughly by inversion. Do not vortex! Invert the tube an additional 5 times after the sample turns completely yellow. *The sample will turn yellow when the neutralization is complete, and a yellowish precipitate will form.*
4. Ensure the plug is attached to the Luer Lock at the bottom of the **ZymoPURE™ Syringe Filter-X**. Place the syringe filter upright in a tube rack and load the lysate in the ZymoPURE™ Syringe Filter-X⁴ and wait 5-8 minutes for the precipitate to float to the top.
5. Remove the Luer Lock plug from the bottom of the syringe and place it into a clean 50 ml conical tube. Place the plunger in the syringe and push the solution through the ZymoPURE™ Syringe Filter-X in one continuous motion until approximately 35 ml of cleared lysate is recovered. Save the cleared lysate!

The volume of lysate recovered from the syringe filter is critical for optimal plasmid DNA binding to the spin-column. Please refer to page 10 in the appendix regarding the adjustment of the volume of ZymoPURE™ Binding Buffer used in step 6 if the clarified lysate volume is below approximately 35 ml.

6. Add 14 ml of **ZymoPURE™ Binding Buffer** to the cleared lysate from step 5 and mix thoroughly by inverting the tube 8 times.

To continue processing the lysate using the recommended vacuum protocol, proceed to page 6. If a vacuum is not available, proceed to page 7 for an alternative centrifugation method.

¹ Lytic enzymes other than lysozyme will require optimization and validation in the **ZymoPURE™ P1** buffer prior to use.

² Incubation times will vary depending on the culture volume, cell density, and age of cells. Harvesting the cells at early log phase is recommended for optimal cell wall digestion.

³ Do not allow the lysis reaction to proceed for more than 3 minutes. Excessive lysis can result in denatured plasmid DNA.

⁴ If the precipitate has formed a homogenous layer at the surface of the neutralized lysate, then invert the tube 3-4 times prior to loading the lysate into the **ZymoPURE™ Syringe Filter-X**.

Adjusting Volume of Binding Buffer

The ratio of lysate to binding buffer is critical for optimal plasmid DNA binding to the spin-column. Therefore, it is important that approximately 35 ml of cleared lysate is recovered from the ZymoPURE™ Syringe Filter-X during step 5 of the protocol. If the clarified lysate volume is below approximately 35 ml, please adjust the volume of ZymoPURE™ Binding Buffer used in step 6 of the protocol. This can be accomplished by multiplying the volume of recovered lysate by 0.4. Please see the example and table below for reference.

Example: For 31 ml of cleared lysate, you will add 12.4 ml of ZymoPURE™ Binding Buffer to the cleared lysate instead of 14 ml in step 6 of the protocol ($31 \text{ ml} \times 0.4 = 12.4 \text{ ml}$).

Optimal Volume of ZymoPURE™ Binding Buffer for Various Volumes of Lysate

Approximate Neutralized Lysate Volume	Volume of ZymoPURE Binding Buffer to Add
35 ml	14.0 ml
34 ml	13.6 ml
33 ml	13.2 ml
32 ml	12.8 ml
31 ml	12.4 ml
30 ml	12.0 ml

Troubleshooting

Problem	Possible Causes and Suggested Solutions
	<p>Poor aeration of culture. The optimal culture volume to air volume ratio is 1:5 or less. For best aeration, use baffled culture flasks, or a vented or gas-permeable seal on the culture vessel.</p> <p>The culture was overgrown, undergrown, contaminated, or antibiotics were omitted from the growth medium. Use a fresh culture for optimal performance. An OD₆₀₀ of 0.2-0.35 is the optimal optical density of a tenfold dilution of the culture.</p> <p>Too much culture used. Lysis and neutralization will be incomplete and the ZymoPURE™ Syringe Filter-X may clog during filtration. <u>More culture does not always equal more plasmid.</u> Incomplete lysis and neutralization are two of the most common causes of failed plasmid preps and both are caused by too much culture being used.</p> <p>Incomplete lysis: After addition of ZymoPURE™ P2, the solution should change from opaque pink to a clear viscous purple, indicating complete lysis. Different <i>E. coli</i> strains often require different growth conditions and may vary in their susceptibility to alkaline lysis.</p>
Low DNA Yield	<p>Incomplete neutralization: The solution should not be viscous following neutralization and the yellowish precipitate should appear fluffy and readily float to the surface. Make sure the neutralization is complete prior to centrifugation. Invert the tube an additional 3-4 times after the sample turns yellow following the addition of ZymoPURE™ P3.</p> <p>ZymoPURE™ P2 and/or ZymoPURE™ Binding Buffer may have precipitated during shipping. To completely resuspend the buffers, incubate the bottles at 30-37°C for 10 minutes and mix by inversion. DO NOT MICROWAVE.</p> <p>ZymoPURE™ Wash 2: Ensure that the correct volume of ethanol was added to the ZymoPURE™ Wash 2 prior to use. Also, ensure that the bottle cap is screwed on tightly after each use to prevent evaporation of the ethanol.</p>

Problem	Possible Causes and Suggested Solutions
<p>Low DNA Yield</p>	<p>Less than approximately 35 ml of neutralized lysate was used for the binding step. The ratio of binding buffer to lysate is critical for optimal plasmid DNA binding to the spin-column. Plasmid DNA yield will be reduced if less than approximately 35 ml is recovered from the syringe filter. Please refer to page 10 in the appendix if less than approximately 35 ml of lysate is recovered from the syringe filter.</p> <p>Incomplete elution: For large size plasmids (> 10 kb), add ZymoPURE™ Elution Buffer and incubate the column for 5-10 minutes before centrifugation. Also, pre-warm the ZymoPURE™ Elution Buffer to 50 °C prior to elution.</p>
<p>Low DNA Quality</p>	<p>Incomplete neutralization: Incomplete neutralization generates poor quality supernatant. Ensure that neutralization is complete by inverting the sample an additional 5 times after the sample turns yellow following the addition of ZymoPURE™ P3.</p> <p>Insufficient centrifugation: Make sure that all centrifugation steps are performed at the indicated speed and time. If a lower centrifuge speed is used, then extend the centrifugation time to compensate.</p>
<p>RNA in eluate</p>	<p>Ensure that ZymoPURE™ P1 has been stored at 4°C. RNase A can be purchased separately if necessary.</p>
<p>Genomic DNA in eluate</p>	<p>Improper handling: Sample was vortexed or handled too roughly. Genomic DNA contamination is usually caused by excessive mechanical shearing during the lysis and neutralization steps. Also, incomplete lysis or neutralization may contribute to genomic DNA contamination in your eluate.</p> <p>Overgrown culture. Overgrown or old cultures may contain more genomic DNA contamination than fresh cultures.</p>

Ordering Information

Product Description	Catalog No.	Size
ZymoPURE™ II Plasmid Maxiprep Kit	D4202	10 Preps.
	D4203	20 Preps.

Individual Kit Components	Catalog No.	Amount
ZymoPURE™ P1 (Red)	D4200-1-150	150 ml
	D4200-1-210	210 ml
	D4200-1-410	410 ml
ZymoPURE™ P2 (Green)	D4200-2-150	150 ml
	D4200-2-210	210 ml
	D4200-2-410	410 ml
ZymoPURE™ P3 (Yellow)	D4200-3-150	150 ml
	D4200-3-210	210 ml
	D4200-3-410	410 ml
ZymoPURE™ Binding Buffer	D4200-4-150	150 ml
	D4200-4-210	210 ml
	D4200-4-410	410 ml
ZymoPURE™ Wash 1	D4200-5-55	55 ml
ZymoPURE™ Wash 2 (Concentrate)	D4200-6-23	23 ml
ZymoPURE™ Elution Buffer	D4200-7-6	6 ml
	D4200-7-12	12 ml
	D4200-7-30	30 ml
Zymo-Spin™ V-PX Column Assembly w/ 15 ml Reservoir-X and 50 ml Reservoir	C1082-5	5
15 ml Reservoir-X	C1084-25	25
50 ml Reservoir	C1032-25	25
ZymoPURE™ Syringe Filter-X	C1092-5	5
ZymoPURE™ Syringe Plunger	C1037-5	5
EndoZero™ Spin-Columns	C1051-10	10
Collection Tubes	C1001-50	50
	C1001-500	500
	C1001-1000	1000

Complete Your Cloning Workflow

✓ Transfection-grade plasmid DNA from a miniprep

ZymoPURE™ Plasmid Miniprep	Size	Catalog No.
ZymoPURE™ Plasmid Miniprep Kit	10 Preps. 50 Preps. 100 Preps. 400 Preps. 800 Preps.	D4208T D4309 D4210 D4211 D4212

✓ 16 Minute Endotoxin-Free Midi & Maxipreps

ZymoPURE™ II Plasmid Prep Kits	Size	Catalog No.
ZymoPURE™ II Plasmid Midiprep Kit	25 Preps. 50 Preps.	D4200 D4201
ZymoPURE™ II Plasmid Maxiprep Kit	10 Preps. 20 Preps.	D4202 D4203
ZymoPURE™ II Plasmid Gigaprep Kit	5 Preps.	D4204

✓ Simple 20 second High Efficiency Transformations

Mix & Go! Competent Cells	Size	Catalog No.
DH5α	10 x 100 µl aliquots 96 x 50 µl aliquots 96 x 50 µl aliquots PCR Plate	T3007 T3009 T3010
JM109	10 x 100 µl aliquots 96 x 50 µl aliquots	T3019 T3020
Zymo10B	10 x 100 µl aliquots 96 x 50 µl aliquots	T3003 T3005
HB101	10 x 100 µl aliquots 96 x 50 µl aliquots	T3011 T3013
TG1	10 x 100 µl aliquots	T3017

✓ Recover ultra-pure highly concentrated DNA from PCR & other sources

DNA Clean & Concentrator™	Size	Catalog No.
DNA Clean & Concentrator™-5	50 Preps. 200 Preps.	D4003 D4004
ZR-96 DNA Clean-Up Kit™	2 x 96 Preps. 4 x 96 Preps.	D4017 D4018

✓ Rapid extraction of ultra-pure DNA from agarose gels

Zymoclean Gel DNA Recovery™	Size	Catalog No.
Zymoclean™ Gel DNA Recovery Kit	50 Preps. 200 Preps.	D4001 D4002



**100% satisfaction guarantee on all Zymo Research products,
or your money back.**

Zymo Research is committed to simplifying your research with quality products and services. If you are dissatisfied with this product for any reason, please call 1(888) 882-9682.

Integrity of kit components is guaranteed for up to one year from date of purchase. Reagents are routinely tested on a lot-to-lot basis to ensure they provide the highest performance and reliability.

This product is for research use only and should only be used by trained professionals. It is not for use in diagnostic procedures. Some reagents included with this kit are irritants. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

™ Trademarks of Zymo Research Corporation

Several ZymoPURE™ product technologies are subject to U.S. and foreign patents or are patent pending.

pGL3™ is a registered trademark of Promega Corporation.



ZYMO RESEARCH

The **BEAUTY** of **SCIENCE** is to Make Things **SIMPLE**®



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682